

ÉTALE COVERS OF KAWAMATA LOG TERMINAL SPACES AND THEIR SMOOTH LOCI

DANIEL GREB, STEFAN KEBEKUS, AND THOMAS PETERNELL

CONTENTS

1. Introduction	1
2. Facts used in the proof	3
3. Proof in case where the branch locus is finite	4
4. Proof in case where the branch locus is relatively finite	5
5. Proof at general points of the branch locus	8
6. Proof of Theorem 1.1 and Corollaries 1.3 and 1.5	10
References	12

1. INTRODUCTION

Working with a singular variety X , one is often interested in comparing the set of étale covers of X with that of its smooth locus X_{reg} . More precisely, one may ask: “What are the obstructions to extend finite étale covers of X_{reg} to all of X ? How do the étale fundamental groups $\hat{\pi}_1(X)$ and $\hat{\pi}_1(X_{\text{reg}})$ compare?”

For projective varieties with Kawamata log terminal (klt) singularities we answer these questions in part. The main result of this paper, Theorem 1.1, asserts that there are no infinite towers of finite morphisms over a klt base variety where all morphisms are étale in codimension one, but branched over a small set. In a certain sense, this result can be seen as saying that the difference between the étale covers of X and those of X_{reg} is finite if X is klt. As an immediate application, we construct in Corollary 1.3 a finite covering $\tilde{X} \rightarrow X$, étale in codimension one, such that the étale fundamental groups $\hat{\pi}_1(\tilde{X})$ and $\hat{\pi}_1(\tilde{X}_{\text{reg}})$ agree.

Theorem 1.1 (Étale covers of klt spaces and their smooth loci). *Let X be a normal, complex, quasi-projective variety. Assume that there exists a $\mathbb{Q}$ -Weil divisor Δ such that (X, Δ) is Kawamata log terminal (klt). Given a sequence of finite morphisms between normal varieties, étale in codimension one,*

$$(1.1.1) \quad X = X_0 \xleftarrow{\gamma_1} X_1 \xleftarrow{\gamma_2} X_2 \xleftarrow{\gamma_3} X_3 \xleftarrow{\gamma_4} \cdots,$$

then all but finitely many of the morphisms γ_i are étale.

Remark 1.2 (Purity of branch locus). By purity of the branch locus, the assumptions that almost all morphisms γ_i of (1.1.1) are étale in codimension one can also be formulated in one the following, equivalent ways.

Date: February 8, 2013.

2010 Mathematics Subject Classification. 14J17, 14B05, 14E30, 14B25.

Key words and phrases. Minimal Model Program, Algebraic Fundamental Group, KLT Singularities.

All three authors were supported in part by the DFG-Forschergruppe 790 “Classification of Algebraic Surfaces and Compact Complex Manifolds”. The first named author gratefully acknowledges the support of the Baden–Württemberg–Stiftung through the “Eliteprogramm für Postdoktorandinnen und Postdoktoranden” and by the Institute of Mathematics at Albert-Ludwigs-Universität Freiburg.

(1.2.1) All X_i are étale over the smooth locus of X .

(1.2.2) The morphisms γ_i are étale over the smooth locus of X_{i-1} .

As an immediate consequence, we state:

Corollary 1.3 (Extension of étale covers from the smooth locus of klt spaces). *Let X be a normal, complex, quasi-projective variety. Assume that there exists a $\mathbb{Q}$ -Weil divisor Δ such that (X, Δ) is klt. Then, there exists a normal variety $\tilde{X}$ and a finite morphism $\gamma : \tilde{X} \rightarrow X$, étale in codimension one, such that the following holds*

(1.3.1) *Any finite étale cover of $\tilde{X}_{\text{reg}}$ extends to a finite étale cover of $\tilde{X}$.*

(1.3.2) *The natural map $\hat{\iota}_* : \hat{\pi}_1(\tilde{X}_{\text{reg}}) \rightarrow \hat{\pi}_1(\tilde{X})$ of étale fundamental groups induced by the inclusion of the smooth locus, $\iota : \tilde{X}_{\text{reg}} \rightarrow \tilde{X}$, is an isomorphism.*

In fact, a somewhat more general statement is true, cf. Remark 6.2. To avoid any potential for confusion, we briefly recall the definitions used in the formulation of Corollary 1.3.

Remark 1.4 (Étale fundamental group). Denoting the complex spaces associated with $\tilde{X}$ and $\tilde{X}_{\text{reg}}$ by $\tilde{X}^{\text{an}}$ and $\tilde{X}_{\text{reg}}^{\text{an}}$, the étale fundamental groups $\hat{\pi}_1(\tilde{X}^{\text{an}})$ and $\hat{\pi}_1(\tilde{X}_{\text{reg}}^{\text{an}})$ are the profinite completions of the usual topological fundamental groups $\pi_1(X^{\text{an}})$ and $\pi_1(X_{\text{reg}}^{\text{an}})$.

As a further application of Theorem 1.1, we obtain a slight generalisation of a recent result by Chenyang Xu, [Xu12, Thm. 2].

Corollary 1.5. *Let X be a normal, complex, projective variety. Assume that there exists a $\mathbb{Q}$ -Weil divisor Δ such that (X, Δ) is klt, and $-(K_X + \Delta)$ is big and nef. Then, the étale fundamental group $\hat{\pi}_1(X_{\text{reg}})$ is finite.*

Further applications will appear in a forthcoming, extended version of this paper.

1.1. Outline of the paper. For the reader's convenience, we have gathered in Section 2 a number of facts which will later be used in the proofs. Most of these are known to experts, but often a little hard to find in the literature.

Starting from a recent result of Chenyang Xu [Xu12], Sections 3–5 discuss and prove variants of Theorem 1.1 in a number of special cases, with increasing order of complexity. With these preparations at hand, Theorem 1.1 and Corollaries 1.3, 1.5 will be shown in the concluding Section 6.

1.2. Acknowledgements. We would like to thank Patrick Graf, Annette Huber-Klawitter and Matthias Wendt for numerous discussions. Chenyang Xu kindly answered our questions by e-mail.

1.3. Notation and global conventions. Throughout the present paper, all schemes, varieties and morphisms will be defined over the complex number field. We follow the notation and conventions of Hartshorne's book [Har77]. In particular, varieties are always assumed to be irreducible.

In the course of the proof, we frequently need to switch between the Zariski- and the Euclidean topology. We will consistently use the following notation.

Notation 1.6 (Complex space associated with a variety). Given a variety or projective scheme X , denote the associated complex space by X^{an} . It is important that X^{an} is endowed with the Euclidean topology, not the Zariski topology. If $f : X \rightarrow Y$ is any morphism of varieties or schemes, denote the induced map of complex spaces by $f^{\text{an}} : X^{\text{an}} \rightarrow Y^{\text{an}}$.

The following trivial convention will later help to formulate inductive arguments concisely. For brevity of notation, we briefly recall the definition of a quasi-étale morphism.

Notation 1.7 (Dimension of the empty set). The empty set has dimension minus infinity, $\dim \emptyset = -\infty$.

Definition 1.8 (Quasi-étale morphisms). A morphism $f : X \rightarrow Y$ between normal schemes of finite type is called quasi-étale if f is of relative dimension zero and étale in codimension one.

2. FACTS USED IN THE PROOF

2.1. Topological triviality of algebraic morphisms. The proof of our main theorem relies on the fact that any the morphism between complex spaces $f^{an} : X^{an} \rightarrow Y^{an}$ which comes from an algebraic morphism $f : X \rightarrow Y$ of complex schemes, has the structure of a topologically trivial fibre bundle, at least over a suitable Zariski-dense subset of the base. To be more precise, we will use the following result.

Proposition 2.1 (Topological triviality of morphisms between pointed varieties). Let $\phi : X \rightarrow B$ be a morphism between normal quasi-projective complex varieties. Let $S \subset X$ be a closed subvariety such that $\phi|_S : S \rightarrow B$ is finite and étale. Then, there exists a Zariski-open, dense subset $U \subseteq B$ with preimage $X_U \subset X$ such that the restriction of ϕ^{an} to X_U^{an} is a topologically locally trivial fibre bundle of pointed spaces with their Euclidean topology.

More precisely, given any point $b \in U$ with fibres $X_b \subset X$, $S_b \subset S$, there exists a neighbourhood $V = V(p) \subseteq U^{an}$, open in the Euclidean topology, with preimage $X_V^{an} := (\phi^{an})^{-1}(V)$, and a commutative diagram of continuous maps,

$$\begin{array}{ccccc} X_b^{an} \times V & \xrightarrow[\text{homeomorphic}]{\varphi_V} & X_V^{an} & \xrightarrow{\text{inclusion}} & X^{an} \\ \text{projection} \downarrow & & \downarrow \phi^{an}|_{X_V^{an}} & & \downarrow \phi^{an} \\ V & \xrightarrow{=} & V & \xrightarrow{\text{inclusion}} & B^{an}, \end{array}$$

such that

$$(2.1.1) \quad \varphi_V^{-1}(S^{an} \cap X_V) = S_b^{an} \times V.$$

Proposition 2.1 is known to experts, but does not appear in the literature in the exact form given above. For the reader's convenience, we sketch a brief proof, reducing Proposition 2.1 to a classical result of Verdier.

Sketch of proof. Except for Equation (2.1.1), the statement of Proposition 2.1 is contained in the work of Verdier, [Ver76, Cor. 5.1], which also holds in the non-normal setting. Proposition 2.1 can easily be deduced by applying Verdier's result to the non-normal, reducible complex scheme

$$X' := (X \times \{0\}) \cup (S \times \mathbb{A}^1) \subset X \times \mathbb{A}^1$$

and to the induced map $f' : X' \rightarrow B$, $(x, z) \mapsto f(x)$. $\square$

For an alternative proof of Proposition 2.1, apply [Kol95, Lem. 2.8.1] to a relative Nagata compactification of $f : X \rightarrow B$.

2.2. Normal spaces, coverings. We will later apply the following connectivity statement to Euclidean-open subsets of normal varieties. Its proof is fairly standard, cf. the proof of the “topological form” of Zariski’s Main Theorem in [Mum88, Chap. III, § 9].

Proposition 2.2 (Connectivity). *Let X be a normal connected complex space. If $S \subset X$ is any proper analytic subset, then $X \setminus S$ is connected.* $\square$

To prove that the morphisms γ_i of Theorem 1.1 are étale, we will use the following characterisation theorem from [Gro71].

Proposition 2.3 (Characterisation of étale morphisms). *Let $f : X \rightarrow Y$ be a finite morphism of reduced quasi-projective schemes. Assume that Y is normal. Then, f is étale if and only if the function Φ that associates to each point $y \in Y$ the number of points in the preimage,*

$$\Phi : \{\text{closed points of } Y\} \rightarrow \mathbb{N}, \quad y \mapsto \#\{\text{closed points of } f^{-1}(y)\},$$

is constant on Y .

Proof. Since étalité is local on Y , we can assume without loss of generality that Y is irreducible. Since f is finite, note that each of the two assumptions “ f étale” and “ Φ constant” implies that every irreducible component of X surjects onto Y . The statement thus reduces to [Gro71, Chapt. I, Thm. 10.11]. $\square$

Finally, we will need the following elementary criterion for a covering space to be homeomorphic to a product.

Proposition 2.4 (Covering spaces of products). *Consider a product $A \times B$ of two connected locally path-connected topological spaces, where B is contractible. Let $\gamma : X \rightarrow A \times B$ be a continuous map which is finite and unbranched. Given any point $b \in B$, write $X_b := \gamma^{-1}(A \times \{b\})$. Then there exists a commutative diagram of continuous maps,*

$$\begin{array}{ccc} X_b \times B & \xrightarrow{\text{homeomorphic}} & X \\ \downarrow (\text{projection}_A \circ \gamma) \times \text{Id}_B & & \downarrow \gamma \\ A \times B & \xlongequal{\quad} & A \times B. \end{array}$$

$\square$

3. PROOF IN CASE WHERE THE BRANCH LOCUS IS FINITE

For varieties with isolated singularities, Theorem 1.1 has already been shown by Chenyang Xu. The following slight generalisation is a direct consequence of his work.

Proposition 3.1 (Main theorem in case where the branch locus is finite). *Let X be a normal quasi-projective variety of dimension $\dim X \geq 2$. Assume that there exists a $\mathbb{Q}$ -Weil divisor Δ such that (X, Δ) is klt. Let*

$$X = X_0 \xleftarrow{\gamma_1} X_1 \xleftarrow{\gamma_2} X_2 \xleftarrow{\gamma_3} X_3 \xleftarrow{\gamma_4} \cdots,$$

be a sequence of finite morphisms between normal varieties, and let $S \subset X$ be a finite set such that for any number $i \in \mathbb{N}^+$ the morphism γ_i is étale away from the finite preimage set $(\gamma_i \circ \cdots \circ \gamma_1)^{-1}(S)$. Then, all but finitely many of the morphisms γ_i are étale.

Proof. Proposition 3.1 is an immediate consequence of [Xu12, Thm. 1], which asserts that for each of the finitely many points $p \in S$, the algebraic local fundamental group,

$$\begin{aligned}\hat{\pi}_1^{\text{loc}}(X, p) &= \text{profinite completion of } \pi_1^{\text{loc}}(X, p) \\ &= \text{profinite completion of } \pi_1(\text{Link}(X, p))\end{aligned}$$

is finite. $\square$

4. PROOF IN CASE WHERE THE BRANCH LOCUS IS RELATIVELY FINITE

The following proposition is a major step towards the proof of Theorem 1.1. Proposition 4.1 handles the case where X is fibred, and where almost all of the morphisms γ_i are known to be étale away from a closed subset S which maps finitely onto the base.

Proposition 4.1 (Main theorem in case where the branch locus is relatively finite). *Let X be a normal quasi-projective variety. Assume that there exists a $\mathbb{Q}$ -Weil divisor Δ such that (X, Δ) is klt, assume that we are given morphisms between normal quasi-projective varieties,*

$$\begin{array}{c} X = X_0 \xleftarrow{\gamma_1} X_1 \xleftarrow{\gamma_2} X_2 \xleftarrow{\gamma_3} X_3 \xleftarrow{\gamma_4} \dots \\ \downarrow \phi \\ B \end{array},$$

and a closed reduced subscheme $S \subset X$ of codimension $\text{codim}_X S \geq 2$, satisfying the following additional conditions.

- (4.1.1) The variety B is smooth.
- (4.1.2) For all $b \in B$, the fibre $X_b := \phi^{-1}(b)$ is reduced, normal, and none of its components are contained in the support of Δ . The cycle $\Delta_b := \Delta \cap X_b$ is a $\mathbb{Q}$ -Weil divisor on X_b . The pair (X_b, Δ_b) is klt.
- (4.1.3) The restriction $\phi|_S : S \rightarrow B$ is finite, étale. In particular, S is smooth.
- (4.1.4) The induced morphism of complex spaces, $\phi^{\text{an}} : X^{\text{an}} \rightarrow B^{\text{an}}$, is a topologically locally trivial fibre bundle of pointed spaces with their Euclidean topology. In other words, the conclusion of Proposition 2.1 holds without shrinking B .
- (4.1.5) For each number $i \in \mathbb{N}^+$, the morphism γ_i is finite. It is étale away from the preimage set $(\gamma_1 \circ \dots \circ \gamma_i)^{-1}(S)$.

Then, all but finitely many of the morphisms γ_i are étale.

The proof of Proposition 4.1 makes use of the following lemma.

Lemma 4.2. *In the setup of Proposition 4.1, assume that S is not empty. Given any fixed index $i \in \mathbb{N}^+$, write $\gamma := \gamma_1 \circ \dots \circ \gamma_i : X_i \rightarrow X$ and consider the reduced preimage $T := (\gamma^{-1}(S))_{\text{red}}$. Then, the induced finite morphism $\gamma|_T : T \rightarrow S$ is étale.*

Remark 4.3. In the setup of Lemma 4.2, given any index $i \in \mathbb{N}$, consider the reduced preimage of S ,

$$S_i := \begin{cases} (\gamma^{-1}(S))_{\text{red}} & \text{if } i > 0 \\ S & \text{if } i = 0 \end{cases}$$

As an immediate consequence of Lemma 4.2, we obtain that the restricted morphism $\gamma|_{S_i} : S_i \rightarrow S_{i-1}$ is étale, cf. [Mil80, Cor. 3.6].

Lemma 4.2 will be shown in the subsequent Section 4.1. The proof of Proposition 4.1 is given in Section 4.2, starting from Page 7.

4.1. Proof of Lemma 4.2. Since γ is finite, its restriction $g := \gamma|_T$ is clearly a finite map. As S is smooth, hence normal, recall from Proposition 2.3 that étalité of g will be established as soon as we show that the function

$$\{\text{closed points of } S\} \rightarrow \mathbb{N}^+, \quad p \mapsto \#\{\text{closed points of } g^{-1}(p)\}$$

is locally constant. We argue by contradiction and assume to the contrary. By semicontinuity, [Gro71, Chapt. I, Thm. 10.11], this amounts to the following.

Assumption 4.4. *There exists a closed point $p \in S$, a Zariski-open subset $S^\circ \subset S$ and a number $d \in \mathbb{N}^+$ such that the following holds.*

- *The point p is contained in the closure of S° .*
- *For all closed points $q \in S^\circ$, the number of closed points in the preimage $g^{-1}(q)$ equals d .*
- *The number of closed points in the preimage of $g^{-1}(p)$ is less than d .*

The fibre bundle structure. We will use the assumption that ϕ^{an} is a topologically locally trivial fibre bundle. To set up notation, set $b := \phi(p) \in B$. Assumption (4.1.4) allows to find connected Euclidean open neighbourhoods $U = U(b) \subset B^{an}$ and $V = V(p) \subset X^{an}$ such that the following holds. For all points $a \in U$, there exists a commutative diagram of continuous maps,

$$\begin{array}{ccc} V_a \times U & \xrightarrow[\text{homeomorphic}]{\varphi} & V \\ \text{projection} \downarrow & & \downarrow \phi^{an}|_V \\ U & \xrightarrow{=} & U \end{array} \quad \text{where} \quad V_a := (\phi^{an})^{-1}(a) \cap V$$

such that

$$(4.5.1) \quad \varphi^{-1}(S^{an}) = (V_a \cap S^{an}) \times U.$$

Simplifying assumptions. Everything said until now remains true when we replace U by a smaller neighbourhood of b and shrink V accordingly. This allows to make the following assumption, which will later simplify the arguments substantially.

Assumption w.l.o.g. 4.6. *The set U is contractible.*

Let $S_p \subset S$ be the connected component of S which contains p . It follows immediately from Assumption 4.4 that there exist points $a \in U$ such that

$$\#\{\text{closed points in } g^{-1}(q)\} = d \quad \text{for any closed point } q \in \phi^{-1}(a) \cap S_p.$$

Choose one such a and fix that choice for the remainder of the proof. Shrinking V , we can then assume that the following additional properties hold.

Assumption w.l.o.g. 4.7. *There exists a unique point $q \in V_a \cap S^{an}$.*

Assumption w.l.o.g. 4.8. *The set $(\gamma^{an})^{-1}(V_a)$ has exactly d distinct connected components.*

End of proof. Consider the preimage $W := (\gamma^{an})^{-1}(V) \subset X_i^{an}$. In order to arrive at a contradiction, we analyse the covering map $\gamma|_{W \setminus T^{an}} : W \setminus T^{an} \rightarrow V \setminus S^{an}$, which is unbranched by Assumption (4.1.4). We have homeomorphisms

$$\begin{aligned} V \setminus S^{an} &\cong (V_a \setminus S^{an}) \times U && \text{Equation (4.5.1)} \\ &\cong (V_a \setminus \{q\}) \times U && \text{Assumption 4.7.} \end{aligned}$$

Since U is contractible by Assumption 4.6, the description of covering spaces given in Proposition 2.4 yields homeomorphisms,

$$W \setminus T^{an} = (\gamma^{an})^{-1}(V \setminus S^{an}) \cong \left((\gamma^{an})^{-1}(V_a \setminus \{q\}) \right) \times U.$$

In particular, Assumption 4.8 implies the following.

Consequence 4.9. *The set $W \setminus T^{an}$ has at least d connected components.*

On the other hand, the morphism $\gamma^{an}|_W : W \rightarrow V$ is finite, so that the preimage set $(\gamma^{an})^{-1}(p) = (g^{an})^{-1}(p)$ intersects every connected component of W . Assumption 4.4 therefore implies the following.

Consequence 4.10. *The set W has strictly less than d connected components.*

Consequences 4.9 and 4.10 contradict the normality of X_i and X_i^{an} . As we have recalled in Proposition 2.2, the connected components of the normal complex space $W \subset X_i$ cannot become disconnected by removing the proper analytic subset $T^{an} \cap W$. This contradiction finishes the proof of Lemma 4.2. $\square$

4.2. Proof of Proposition 4.1. We aim to apply Proposition 3.1 to a very general fibre of the morphism ϕ . Once it is shown that almost all of the morphisms γ_i are étale along this one fibre, we will employ Lemma 4.2 to conclude that they are in fact everywhere étale.

Setup and notation. If S is empty, there is nothing to show. We will therefore assume throughout the proof that $S \neq \emptyset$. Given any index $i \in \mathbb{N}$, consider the map

$$\Gamma_i := \begin{cases} (\gamma_1 \circ \cdots \circ \gamma_i) : X_i \rightarrow X. & \text{if } i > 0 \\ \text{Id} : X_0 \rightarrow X & \text{if } i = 0 \end{cases}$$

and the reduced preimage subscheme

$$S_i := ((\Gamma_i)^{-1}(S))_{\text{red}} \subset X_i.$$

By Seidenberg's theorem, [BS95, Thm. 1.7.1], there exists a dense open subset $B_i^\circ \subseteq B$ such that the following holds for any closed point $b \in B_i^\circ$.

(4.11.1) The scheme-theoretic fibre $X_{i,b} := (\phi \circ \Gamma_i)^{-1}(b)$ is reduced and normal.

Choice of a good base point. Owing to the uncountability of our base field $\mathbb{C}$, the intersection of the countably many dense, Zariski-open sets $(B_i^\circ)_{i \in \mathbb{N}}$ is not empty. We can therefore find (uncountably many) closed points $b \in B$ such that Property (4.11.1) holds for all indices $i \in \mathbb{N}$. Choose one such b and fix that choice for the remainder of the proof.

Induced morphisms between the fibres. Consider the restricted morphisms $\gamma_{i,b} := \gamma_i|_{X_{i,b}} : X_{i,b} \rightarrow X_{i-1,b}$. The $\gamma_{i,b}$ align to give a sequence of finite morphisms,

$$(4.11.2) \quad X_{0,b} \xleftarrow{\gamma_{1,b}} X_{1,b} \xleftarrow{\gamma_{2,b}} X_{2,b} \xleftarrow{\gamma_{3,b}} X_{3,b} \xleftarrow{\gamma_{4,b}} \cdots$$

Recalling from Assumption (4.1.5) that the morphisms γ_i are étale away from the preimage set S_i , it follows from stability of étalité under base change, [Mil80, Prop. 3.3], that the restricted morphisms $\gamma_{i,b}$ are étale away from the finite set

$$X_{i,b} \cap S_i = (\gamma_{1,b} \circ \cdots \circ \gamma_{i,b})^{-1}(S_0 \cap X_{0,b}).$$

Together with Assumption (4.1.2) and Property (4.11.1), this allows to apply Proposition 3.1 to the sequence of morphisms given in (4.11.2). We obtain that all but finitely many of the morphisms $\gamma_{i,b}$ are étale.

End of proof. Let $i \in \mathbb{N}^+$ be one of the indices for which $\gamma_{i,b}$ is étale. We will then show that γ_i is étale. Using the characterisation of étalité given in Proposition 2.3, it suffices that show that

$$(4.11.3) \quad \#\{\text{closed points of } \gamma_i^{-1}(p)\} = \deg \gamma_i \quad \text{for all } p \in S_{i-1}.$$

Lemma 4.2 and Remark 4.3 simplify this task: since $\gamma_i|_{S_i} : S_i \rightarrow S_{i-1}$ is étale, it suffices to check Equation (4.11.3) for finitely many closed points $p \in S_{i-1}$, one for each connected component of S_{i-1} . By Assumption (4.1.3), the map $(\phi \circ \Gamma_{i-1})|_{S_{i-1}} : S_{i-1} \rightarrow B$ is finite, and every connected component of S_{i-1} will intersect $X_{i-1,b}$. It will therefore suffice to check Equation (4.11.3) for the finitely many closed points of $X_{i-1,b} \cap S_{i-1}$.

By assumption, $\gamma_{i,b}$ is étale and the function

$$\{\text{closed points of } X_{i-1,b}\} \rightarrow \mathbb{N}^+, \quad p \mapsto \#\{\text{closed points of } \gamma_i^{-1}(p)\}$$

therefore locally constant. It takes the value $\deg \gamma_i$ on the dense open set $X_{i-1,b} \setminus S_{i-1}$, over which γ_i is assumed to be étale by (4.1.5). Equation (4.11.3) therefore holds true for all closed points of $X_{i-1,b}$, including those contained in S_{i-1} . This finishes the proof of Proposition 4.1. $\square$

5. PROOF AT GENERAL POINTS OF THE BRANCH LOCUS

As a next step in the proof of Theorem 1.1 we show that almost all of the morphisms γ_i are étale over general points of any potential branch locus.

Proposition 5.1. *Let X be a normal quasi-projective variety. Assume that there exists a $\mathbb{Q}$ -Weil divisor Δ such that (X, Δ) is klt. Assume further that we are given a sequence of finite morphisms between normal varieties,*

$$X = X_0 \xleftarrow{\gamma_1} X_1 \xleftarrow{\gamma_2} X_2 \xleftarrow{\gamma_3} X_3 \xleftarrow{\gamma_4} \dots,$$

and a closed reduced subscheme $S \subset X$ of codimension $\text{codim}_X S \geq 2$ such that for any number $i \in \mathbb{N}^+$, the morphism γ_i is étale away from the reduced preimage

$$S_i := ((\gamma_1 \circ \dots \circ \gamma_i)^{-1}(S))_{\text{red}}.$$

If $T \subseteq S$ is any irreducible component, then there exists a dense, open subscheme $T^\circ \subset T$ and a number $N_T \in \mathbb{N}^+$ such that for all $i \geq N_T$, the morphism γ_i is étale at all points of the reduced preimage

$$T_i^\circ := ((\gamma_1 \circ \dots \circ \gamma_i)^{-1}(T^\circ))_{\text{red}}.$$

5.1. Proof of Proposition 5.1. If S is empty, there is nothing to show. If T is an isolated point of S , the assertion has already been shown in Proposition 3.1. We will therefore make the following assumption throughout the proof.

Assumption w.l.o.g. 5.2. *The variety T is positive-dimensional.*

Projection to the subvariety T . We will analyse the behaviour of the morphisms γ_i near the generic point of T using the technique of “projection to a subvariety”, explained in detail in [GKKP11, Sect. 2.G]. More specifically, recall from [GKKP11, Prop. 2.26] that there exists a Zariski-open subset $X^\circ \subseteq X$ such that $T^\circ := T \cap X^\circ$ is not empty, and normal varieties Z, B forming a diagram

$$\begin{array}{ccc} Z & \xrightarrow[\text{finite, étale}]{\psi} & X^\circ \\ \phi \downarrow & & \\ B & & \end{array}$$

with the property that the restriction of ϕ to any connected component of $\psi^{-1}(T^\circ)$ is an isomorphism. In particular, $\phi|_{\psi^{-1}(T^\circ)} : \psi^{-1}(T^\circ) \rightarrow B$ is étale. Given any number $i \in \mathbb{N}$, let Z_i be the normalisation of the fibre product $Z \times_X X_i$. These spaces and their standard projections form a commutative ladder,

$$(5.2.1) \quad \begin{array}{ccccccc} X = X_0 & \xleftarrow{\gamma_1} & X_1 & \xleftarrow{\gamma_2} & \cdots & \xleftarrow{\gamma_{N^{k+1}}} & X_{N^{k+1}} & \xleftarrow{\cdots} \\ \psi_0 = \psi \uparrow & & \psi_1 \uparrow & & & & \psi_{N^{k+1}} \uparrow & \\ Z = Z_0 & \xleftarrow{\phi_1} & Z_1 & \xleftarrow{\phi_2} & \cdots & \xleftarrow{\phi_{N^{k+1}}} & Z_{N^{k+1}} & \xleftarrow{\cdots} \\ \downarrow \phi & & & & & & & \\ B & & & & & & & \end{array}$$

satisfying the following properties for all $i \in \mathbb{N}^+$.

(5.2.2) All horizontal arrows are finite, and étale away from the proper closed subschemes $S_i \subset X_i$ and $\psi_i^{-1}(S_i) \subset Z_i$, respectively.

(5.2.3) All vertical arrows other than ϕ are étale, mapping the spaces Z_i finitely onto the Zariski-open sets $X_i^\circ := (\gamma_1 \circ \cdots \circ \gamma_i)^{-1}(X^\circ)$.

Application of Proposition 4.1. Shrinking X° if necessary, we are free to assume that the following additional properties hold.

Assumption w.l.o.g. 5.3. *The intersection $X^\circ \cap S$ is contained in T and therefore equal to the open subset $T^\circ \subseteq T$. By stability of étalité under base change, [Mil80, Prop. 3.3], we have that for any number $i \in \mathbb{N}^+$, the morphism ϕ_i is étale away from the reduced preimage*

$$((\phi_1 \circ \cdots \circ \phi_i)^{-1}(\psi^{-1}(T^\circ)))_{\text{red}}.$$

Since ψ is étale, it is possible to pull back Weil divisors from X to Z . Writing $\Delta_Z := \psi^{-1}(\Delta)$, it follows from [KM98, Prop. 5.20] that the pair (Z, Δ_Z) is klt. Shrinking B , Z and X° , using generic smoothness of B , Seidenberg's theorem, [BS95, Thm. 1.7.1], the Bertini theorem for klt pairs, [KM98, Lem. 5.17] as well as the topological triviality of morphisms between varieties discussed in Proposition 2.1, we can additionally assume that the following holds.

Assumption w.l.o.g. 5.4 (Generic Smoothness). *The variety B is smooth.*

Assumption w.l.o.g. 5.5 (Seidenberg and Bertini theorem for klt pairs). *For all $b \in B$, the fibre $Z_b := \phi^{-1}(b)$ is reduced, normal, and none of its components are contained in the support of Δ_Z . The cycle $\Delta_b := \Delta_Z \cap Z_b$ is a $\mathbb{Q}$ -Weil divisor on Z_b . The pair (Z_b, Δ_b) is klt.*

Assumption w.l.o.g. 5.6 (Topological triviality). *The induced morphism of complex spaces, $\phi^{an} : Z^{an} \rightarrow B^{an}$ is a topologically locally trivial fibre bundle of pointed spaces with their Euclidean topology.*

In summary, we have just checked that the morphisms $\phi, \phi_1, \phi_2, \dots$ of Diagram (5.2.1) satisfy all assumptions made in Proposition 4.1. Consequently, all but finitely many of the morphisms $(\phi_i)_{i \in \mathbb{N}}$ are étale. In other words, there exists a number $N_T \in \mathbb{N}^+$ such that the morphisms ϕ_i are étale for all $i \geq N_T$.

End of proof. Commutativity of Diagram (5.2.1) together with Property (5.2.3) thus implies that the restrictions $\gamma_i|_{X_i^\circ} : X_i^\circ \rightarrow X_{i-1}^\circ$ are étale for all $i \geq N_T$, cf. [Mil80, Cor. 3.6]. Assumption 5.3 thus implies that the morphisms γ_i are étale at all points $q \in T_i^\circ$, as claimed. $\square$

6. PROOF OF THEOREM 1.1 AND COROLLARIES 1.3 AND 1.5

6.1. Proof of Theorem 1.1. Theorem 1.1 is an immediate consequence of the following proposition, whose statement is more involved, but amenable to induction. Theorem 1.1 follows from Proposition 6.1 by using Remark 1.2 and taking the singular locus of X for S , that is, $S := X_{\text{sing}}$.

Proposition 6.1. *Let X be a normal quasi-projective variety. Assume that there exists a $\mathbb{Q}$ -Weil divisor Δ such that (X, Δ) is klt. Assume further that we are given a sequence of finite morphisms between normal varieties,*

$$X = X_0 \xleftarrow{\gamma_1} X_1 \xleftarrow{\gamma_2} X_2 \xleftarrow{\gamma_3} X_3 \xleftarrow{\gamma_4} \cdots,$$

and a closed reduced subscheme $S \subset X$ of codimension $\text{codim}_X S \geq 2$ such that for any number $i \in \mathbb{N}^+$, the morphism γ_i is étale away from the reduced preimage

$$S_i := ((\gamma_1 \circ \cdots \circ \gamma_i)^{-1}(S))_{\text{red}}.$$

Then, there exists a number N_S such that for any number $i \geq N_S$, the morphism γ_i is étale.

Proof. We argue by induction, using the dimension of S as our induction variable. If $\dim S = 0$, the claim has already been shown in Proposition 3.1. We will therefore assume that $\dim S > 0$ and that Proposition 6.1 has already been shown for all klt pairs (Y, D) and proper subschemes $R \subset Y$ where $\dim R < \dim S$.

Decompose S into irreducible components, $S = S^1 \cup S^2 \cup \cdots \cup S^n$. Apply Proposition 5.1 to each component S^i and, using the open sets $(S^i)^\circ$ and numbers N_{S^i} introduced in Proposition 5.1, set

$$S' := S \setminus ((S^1)^\circ \cup \cdots \cup (S^n)^\circ) \quad \text{and} \\ N := \max\{N_{S^1}, \dots, N_{S^n}\}.$$

With these definitions, it is clear that for any number $i \geq N$, the morphism γ_i is étale away from the reduced preimage

$$S'_i := ((\gamma_1 \circ \cdots \circ \gamma_i)^{-1}(S'))_{\text{red}},$$

whose dimension is strictly smaller than that of S , that is, $\dim S' < \dim S$.

We aim to apply the induction hypothesis to X_N . As before, using finiteness of $\gamma_1 \circ \cdots \circ \gamma_N$, it is possible to pull the $\mathbb{Q}$ -Weil divisor Δ back to a $\mathbb{Q}$ -Weil divisor Δ_N on X_N . Again, recall from [KM98, Prop. 5.20] that the pair (X_N, Δ_N) is then klt. In summary, we can apply the induction hypothesis to the sequence of finite morphisms

$$X_N \xleftarrow{\gamma_{N+1}} X_{N+1} \xleftarrow{\gamma_{N+2}} X_{N+2} \xleftarrow{\gamma_{N+3}} \cdots,$$

which are unbranched away from the preimages of S'_N . We obtain a number $N_{S'}$ such that the γ_i are étale for all $i \geq N_S := N + N_{S'}$. This finishes the proof of Proposition 6.1. $\square$

6.2. Proof of Corollary 1.3. To prove Corollary 1.3 we will first show that there exists a finite quasi-étale cover $\tilde{X} \rightarrow X$ that satisfies Statement (1.3.1). We will then see in a second step that this cover also satisfies Statement (1.3.2).

Proof of Statement (1.3.1). We argue by contradiction and assume that there is no cover which satisfies Condition (1.3.1). In particular, $\text{Id}_X : X \rightarrow X$ does not satisfy the condition, so that there exists a finite quasi-étale cover $\gamma_1 : X_1 \rightarrow X$ that is not étale. Observing that X_1 also does not satisfy Condition (1.3.1), we can inductively construct a sequence of morphisms,

$$X = X_0 \xleftarrow{\gamma_1} X_1 \xleftarrow{\gamma_2} X_2 \xleftarrow{\gamma_3} X_3 \xleftarrow{\gamma_4} \cdots,$$

where none of the quasi-étale maps is actually étale. This contradiction to Theorem 1.1 finishes the proof of Statement (1.3.1).

Proof of Statement (1.3.2). If $\tilde{X} \rightarrow X$ is any cover for which Statement (1.3.1) holds, injectivity of $\hat{\iota}_*$ follows immediately. For surjectivity, recall from [FL81, 0.7.B on p. 33] and [Kol95, Prop. 2.10] that the push-forward map between topological fundamental groups,

$$\iota_* : \pi_1(\tilde{X}_{\text{reg}}^{\text{an}}) \rightarrow \pi_1(\tilde{X}^{\text{an}})$$

is surjective. Because profinite completion is a right-exact functor, [RZ10, Lem. 3.2.3 and Prop. 3.2.5], it follows that $\hat{\iota}_*$ is likewise surjective. This finishes the proof of Statement (1.3.2) and hence of Corollary 1.3. $\square$

Remark 6.2. In the setting of Corollary 1.3, let $\gamma_1 : X_1 \rightarrow X$ be any finite quasi-étale morphism of normal varieties, and $\Delta_1 := \gamma_1^*(\Delta)$ the pull-back of the divisor Δ , which exists because γ_1 is finite. Recalling from [KM98, Prop. 5.20] that (X_1, Δ_1) is klt, we can apply Corollary 1.3 to X_1 . In summary, we see that any finite quasi-étale cover of X can be enlarged to a cover where Conditions (1.3.1) and (1.3.2) hold.

6.3. Proof of Corollary 1.5. We argue by contradiction and suppose that the étale fundamental group $\hat{\pi}_1(X_{\text{reg}})$ is infinite. We can thus find smooth varieties X_i° and a sequence of étale morphisms,

$$X_{\text{reg}} = X_0^\circ \xleftarrow{\gamma_1^\circ} X_1^\circ \xleftarrow{\gamma_2^\circ} X_2^\circ \xleftarrow{\gamma_3^\circ} X_3^\circ \xleftarrow{\gamma_4^\circ} \cdots,$$

where each of the morphisms γ_i° is given by a non-trivial surjection of $\hat{\pi}_1(X_{i-1}^\circ)$ to a finite group, [RZ10, Thm. 2.6.4]. By a classical result of Grauert-Remmert [DG94, §3.3], [Gro71, XII, 5.4] the morphisms γ_i° admit unique extensions γ_i which a sequence of finite quasi-étale morphisms of normal connected varieties,

$$X = X_0 \xleftarrow{\gamma_1} X_1 \xleftarrow{\gamma_2} X_2 \xleftarrow{\gamma_3} X_3 \xleftarrow{\gamma_4} \cdots.$$

Theorem 1.1 asserts that almost all γ_i are étale. In contrast to that, we will now show that in fact none of the γ_i can possibly be étale.

To this end, use finiteness of γ_i to define pull-back divisors on X_i ,

$$\Delta_i := (\gamma_1 \circ \cdots \circ \gamma_i)^*(\Delta),$$

and use finiteness of the γ_i as well as [KM98, Prop. 5.20] to conclude that the pairs (X_i, Δ_i) are klt. In fact, more is true. Since the γ_i are all étale in codimension one, there exists a $\mathbb{Q}$ -linear equivalence of $\mathbb{Q}$ -Cartier, $\mathbb{Q}$ -Weil divisors,

$$(6.3.1) \quad K_{X_i} + \Delta_i \sim_{\mathbb{Q}} (\gamma_1 \circ \cdots \circ \gamma_i)^*(K_X + \Delta).$$

Equation (6.3.1) shows that the divisors $-(K_{X_i} + \Delta_i)$ are nef and big, for all $i \in \mathbb{N}$. In this setting, a result of Takayama [Tak00, Thm. 1.1], see also [Zha06, Cor. 1], asserts that the associated analytic spaces X_i^{an} are all simply connected. In particular, none of them admits a connected étale cover. This contradiction finishes the proof of Corollary 1.5. $\square$

REFERENCES

- [BS95] Mauro C. Beltrametti and Andrew J. Sommese. *The adjunction theory of complex projective varieties*, volume 16 of *de Gruyter Expositions in Mathematics*. Walter de Gruyter & Co., Berlin, 1995. ↑ 7, 9
- [DG94] Gerd Dethloff and Hans Grauert. Seminormal complex spaces. In *Several complex variables, VII*, volume 74 of *Encyclopaedia Math. Sci.*, pages 183–220. Springer, Berlin, 1994. ↑ 11
- [FL81] William Fulton and Robert Lazarsfeld. Connectivity and its applications in algebraic geometry. In *Algebraic geometry (Chicago, Ill., 1980)*, volume 862 of *Lecture Notes in Math.*, pages 26–92. Springer, Berlin, 1981. ↑ 11
- [GKKP11] Daniel Greb, Stefan Kebekus, Sándor J. Kovács, and Thomas Peternell. Differential forms on log canonical spaces. *Inst. Hautes Études Sci. Publ. Math.*, 114(1):87–169, November 2011. DOI:10.1007/s10240-011-0036-0 An extended version with additional graphics is available as [arXiv:1003.2913](https://arxiv.org/abs/1003.2913). ↑ 8
- [Gro71] Alexandre Grothendieck. *Revêtements étales et groupe fondamental*. Springer-Verlag, Berlin, 1971. Séminaire de Géométrie Algébrique du Bois Marie 1960–1961 (SGA 1), Dirigé par Alexandre Grothendieck. Augmenté de deux exposés de M. Raynaud, Lecture Notes in Mathematics, Vol. 224. ↑ 4, 6, 11
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52. ↑ 2
- [KM98] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. ↑ 9, 10, 11
- [Kol95] János Kollár. *Shafarevich maps and automorphic forms*. M. B. Porter Lectures. Princeton University Press, Princeton, NJ, 1995. ↑ 3, 11
- [Mil80] James S. Milne. *Étale cohomology*, volume 33 of *Princeton Mathematical Series*. Princeton University Press, Princeton, N.J., 1980. ↑ 5, 7, 9
- [Mum88] David Mumford. *The Red Book of Varieties and Schemes*, volume 1358 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1988. ↑ 4
- [RZ10] Luis Ribes and Pavel Zalesskii. *Profinite groups*, volume 40 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 2010. ↑ 11
- [Tak00] Shigeharu Takayama. Simple connectedness of weak Fano varieties. *J. Algebraic Geom.*, 9(2):403–407, 2000. ↑ 11
- [Ver76] Jean-Louis Verdier. Stratifications de Whitney et théorème de Bertini-Sard. *Invent. Math.*, 36:295–312, 1976. ↑ 3
- [Xu12] Chenyang Xu. Finiteness of algebraic fundamental groups. preprint [arXiv:1210.5564](https://arxiv.org/abs/1210.5564), October 2012. ↑ 2, 5
- [Zha06] Qi Zhang. Rational connectedness of log $\mathbb{Q}$ -Fano varieties. *J. Reine Angew. Math.*, 590:131–142, 2006. ↑ 11

DANIEL GREB, ARBEITSGRUPPE ALGEBRA/TOPOLOGIE, FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, 44780 BOCHUM, GERMANY

E-mail address: daniel.greb@rub.de

URL: <http://www.rub.de/ffm/fakultaet/lehrstuehle/index.html>

STEFAN KEBEKUS, MATHEMATISCHES INSTITUT, ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, ECKERSTRASSE 1, 79104 FREIBURG IM BREISGAU, GERMANY

E-mail address: stefan.kebekus@math.uni-freiburg.de

URL: <http://home.mathematik.uni-freiburg.de/kebekus>

THOMAS PETERNELL, MATHEMATISCHES INSTITUT, UNIVERSITÄT BAYREUTH, 95440 BAYREUTH, GERMANY

E-mail address: thomas.peternell@uni-bayreuth.de

URL: <http://www.staff.uni-bayreuth.de/~btm109/peternell/pet.html>